



Changes in the non-crustacean zooplankton community in the middle Adriatic Sea during the Eastern Mediterranean Transient

MIRNA BATISTIĆ
RADE GARIĆ
MIRA MOROVIĆ

¹ University of Dubrovnik, Institute for Marine and Coastal Research, Kneza D. Jude 12, 20000 Dubrovnik, Croatia

² Institute of Oceanography and Fisheries, I. Meštrovića 63, 21000 Split, Croatia

Correspondence:

Mirna Batistić
e-mail: mirna.batistic@unidu.hr

Key words: zooplankton, hydroclimatic changes, Middle Adriatic

Abstract

Background and Purpose: Here we presented changes in the non-crustacean zooplankton community in the years characterized by the large scale changes in the thermohaline circulation in the East Mediterranean known as the Eastern Mediterranean transient (EMT) and stronger inflow of colder and less saline Modified Atlantic Water (MAW) into the Adriatic Sea.

Material and Method: Monthly samplings from February 1995 to February 1996, were performed at fixed station Stončica near the Island of Vis in the open oligotrophic Middle Adriatic waters. Zooplankton samples, taken vertically from 100 m to the surface using a Nansen net with 125- μ m mesh size were analyzed in detail for the following zooplankton taxa: Hydromedusae, Calycophorae, Ctenophora, Pteropoda, Heteropoda, Polychaeta and Chaetognatha.

Results and Conclusions: Among the investigated non-crustacean zooplankton three species were registered for the first time, while one species reappeared after years of absence. Compared with earlier data there was a dramatic change in dominant species of calycophoran medusae, pteropods and polychaetes. Data presented herein provide baseline information that is essential for the evaluation of impact of hydroclimatic changes on the zooplankton community, which started in the East Mediterranean in the 1990s and are still on going.

INTRODUCTION

Adriatic Sea is divided into three parts: the North, the Middle and the South Adriatic. The North Adriatic is very shallow, with an average depth of 30 to 40 m and maximum of 70 m, while the Middle Adriatic is much deeper, reaching 280 m in the Jabuka Pit. It is separated from the deepest South Adriatic by the 170 m deep Palagruža Sill. Adriatic Sea is connected with the Mediterranean Sea through the Otranto Strait. The deepest part of the South Adriatic (cca. 1240 m depth) is influenced by Ionian and eastern Levantine waters, and plays an important role as a site of the formation of dense water, the Adriatic Dense Water (AdDW) (1). The AdDW spreads into the Ionian abyss and represents the main component of the East Mediterranean Deep Water (EMDW).

Furthermore, it has been demonstrated that the Ionian upper layer circulation, the thermohaline properties of the AdDW-EMDW (2, 3) and the salt distribution over the East Mediterranean (4) are interconnected through the BiOS, a feedback mechanism between the South

Received May 26, 2015.
Revised February 18, 2016.
Accepted February 18, 2016.

Adriatic (SA) and Ionian Sea (IS) that changes the upper-layer circulation of the North Ionian Gyre (NIG). During the last 25 years it has been observed that the upper-layer circulation in the Ionian reversed on decadal time scales, from anticyclonic to cyclonic and vice-versa (3, 4, 5). While the cyclonic regime brings warmer and saltier waters of Levantine origin into the Adriatic, the anti-cyclonic pattern favours the inflow of colder and less saline Ionian waters diluted by the Atlantic Water (5). During the early 1990s, the deep water formation areas switched from the SA to the Cretan Sea (6). This event, known as the East Mediterranean Transient (EMT), caused an abrupt change in the East Mediterranean circulation and uniquely strengthened inflow of Modified Atlantic Water (MAW) into the Adriatic (2,7). This interesting phenomenon caused large changes in the Mediterranean marine environment (*e.g.* 4, 5, 8, 9) that are not yet fully known.

BiOS mechanism has important influence on biological and chemical properties of the South Adriatic (3, 5, 11), as well as on physical and chemical properties of the Middle Adriatic (9, 10). The influence of BiOS on plankton communities and the advection of immigrant zooplankton species has only been known for the South Adriatic non-crustacean zooplankton community (11). Magnitude of influence on the zooplankton community in the rest of the Adriatic is unknown to date. We par-

tially addressed this question by analyzing plankton records collected in the Middle Adriatic in 1995/1996, the years characterized by the abrupt change in the eastern Mediterranean circulation (EMT) and advection of Modified Atlantic Water (MAW) in the Adriatic. Zooplankton data originate from the permanent observation site Stončica in the Middle Adriatic which is strongly influenced by incoming water masses from the Ionian Sea. The importance of the open Middle Adriatic in tracing the Adriatic circulation and water masses has been recognized from the beginning of modern oceanographic investigations of the Adriatic (12, 13, 14). Long-term zooplankton data from Stončica have been analysed in several papers (15, 16, 17, 18, 19). However, considering that zooplankton investigations in this area were largely focused on crustacean zooplankton (see in ref. 16), this paper brings valuable additional perspective on changes in biological properties of the water column based on the non-crustacean zooplankton community in the context of large scale influences on thermohaline circulation within this area.

MATERIAL AND METHODS

Monthly samplings from February 1995 to February 1996, were performed at fixed station Stončica near the Island of Vis in the open Middle Adriatic (43°02'38"N 16°17'7"E, depth 106 m, Fig. 1) in the frame of regular




Figure 1. Study area




Figure 2. Temperature and salinity at Stončica station during the investigated period.

monitoring carried out by the Institute of oceanography and fisheries in Split (www.izor.hr). Based on long-term monitoring of the chemical and biological parameters, Stončica site is designated as an oligotrophic open sea site, characterized by high transparency and decreased phytoplankton and zooplankton abundance, in comparison to more productive coastal areas in the Middle Adriatic.

Zooplankton samples, taken vertically from 100 m to the surface using a Nansen net with 125- μ m mesh size (57 cm mouth diameter and 255 cm total length), were preserved in 2.5% formaldehyde and analyzed in detail for the following zooplankton taxa: Hydromedusae, Siphonophorae, Ctenophora, Pteropoda, Heteropoda, Polychaeta and Chaetognatha. Taxonomic identification and

counting of individuals were performed with a Zeiss stereomicroscope at 25x and 40x magnifications. Abundance of all groups except calyctophoran siphonophores is presented as the number of specimens per 10 m³. Calyctophoran abundance was expressed according to the number of nectophores (polygastric stage) of each species (nectophores per 10 m³).

Water samples for oceanographic measurements were taken with 5-litre Niskin bottles at standard depths of 0, 5, 10, 20, 30, 50, 75 and 100 m. Seawater temperature and salinity was measured using automatic CTD probe.

The relationship between groups, dominant species and environmental parameters was tested using the Pearson's rank correlation coefficient. Dataset was log transformed to ensure that the variables normally distributed.

The dominant species within a particular group were determined according to their contribution in total abundance (percentage number, PN) and their occurrence in total number of samples (frequency of occurrence, F) where PN is $\geq 10\%$ and F is $\geq 50\%$ (20, 21, 22, 23, 24). Percentage number was calculated as follows:

$PN = (\pi/P) \times 100$ where π is number of individuals of particular species and P is total number of individuals of all species.

Frequency of occurrence was calculated using the following formula:

$F = (n_i/N) \times 100$ where n_i is number of samples where the species was recorded and N is total number of samples.

RESULTS

Hydrographic parameters

The temperature varied between 13.4°C in January and 23.0°C in July (Fig. 2). Periods of isothermy occurred in January, February, March and November. Between June and September there was a marked thermal stratification,

Table 1. Pearson coefficient of correlation between abundance of zooplankton groups and hydrographic parameters ($n = 11$). Asterisks indicate the level of significance: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Taxa	Temperature	Salinity
Groups		
Hydromedusae	-0.182	-0.288
Calyctophorae	-0.199	-0.322
Heteropoda	0.572*	-0.198
Pteropoda	0.313	-0.706**
Polychaeta	0.128	-0.446
Chaetognatha	0.842***	-0.134

with thermocline positioned between 10 and 20 m depth (June and July), and 30 and 50 m depth (September).

The vertical salinity distribution indicated that major fluctuations occurred in the upper 10 m, with lowest values in July (37.46 at 0 m, Fig. 2). Minimal fluctuations were recorded in winter months and during the autumn isothermal period. Halocline was formed in May and June between 10 and 20 m and in July between 5 and 10 m. Below 20 m, salinity was higher than 38.20 (Fig. 2).

Population structure and abundance of non-crustacean zooplankton

The most abundant groups were Chaetognatha, Pteropoda, Hydromedusae and Calyctophorae (Fig. 3). Annual variations of abundances are presented in Fig. 3. Total abundances of chaetognaths and heteropods were in significant correlation with temperature ($p < 0.001$ and $p < 0.05$, respectively) while the total abundance of pteropods was in correlation with salinity ($p < 0.01$) (Table 1).

During the investigated period 50 species were identified (Table 2): 11 species of Hydromedusae, 8 species of Calyctophorae, 12 species of Pteropoda, 3 species of Het-




Figure 3. Abundances of investigated zooplankton groups.

Table 2. Mean (ind. per 10m³), standard deviation (SD), percentage number (PN, %) and frequency of occurrence (F, %) of the different taxa in the respective group.

Taxa	Mean	SD	PN	F
Hydromedusae				
<i>Sarsia gemmifera</i>	0.07	0.24	0.3	9
<i>Euphysa aurata</i>	0.36	0.89	1.2	18
<i>Laodicea undulata</i>	0.11	0.20	0.4	18
<i>Obelia</i> spp.	0.65	1.24	2.3	27
<i>Clytia hemisphaerica</i>	3.45	5.62	8.7	70
<i>Liriope tetrphylla</i>	5.56	5.23	22.8	100
<i>Aglaura hemistoma</i>	13.35	16.71	46.3	100
<i>Persa incolorata</i>	0.91	1.23	3.1	64
<i>Rhopalonema velatum</i>	1.52	1.08	6.3	70
<i>Solmundella bitentaculata</i>	1.09	1.42	3.8	73
<i>Solmaris leucostyla</i>	0.25	0.41	0.9	30
Calycophorae				
<i>Lensia subtilis</i>	5.85	4.31	16.8	100
<i>Muggiaea atlantica</i>	19.24	27.46	55.3	100
<i>M. kochii</i>	4.64	5.58	12.1	100
<i>Chelophyes appendiculata</i>	0.11	0.26	0.3	18
<i>Eudoxoides spiralis</i>	0.44	0.52	1.3	55
<i>Sphaeronectes gracilis</i>	4.22	4.42	12.1	100
<i>Abylopsis tetragona</i>	0.07	0.24	0.2	9
<i>Bassia bassensis</i>	0.62	1.99	1.8	18
Ctenophora				
<i>Ctenophora</i> unidentified	0.87	1.93		30
Heteropoda				
<i>Atlanta peronii</i>	0.47	1.44	62.0	18
<i>Protatlanta souleyeti</i>	0.25	0.60	33.3	27
<i>Firoloida desmarestia</i>	0.04	0.12	4.8	9
Pteropoda				
<i>Limacina inflata</i>	23.02	44.23	26.2	100
<i>L. trochiformis</i>	3.84	0.69	4.2	70
<i>Styliola subula</i>	0.04	0.12	0.1	9
<i>Creseis virgula</i>	35.85	58.91	43.7	81
<i>C. acicula</i>	14.76	38.79	18.0	54
<i>Hyalocylix striata</i>	0.15	0.37	0.2	18
<i>Clio pyramidata</i>	0.04	0.12	0.1	9
<i>Clio cuspidata</i>	0.04	0.12	0.1	9
<i>Cavolinia inflexa</i>	1.16	3.21	0.3	36
<i>Peracle reticulata</i>	0.47	1.44	0.6	18
<i>Cymbulia peronii</i>	0.47	1.22	0.2	18
<i>Desmopterus papilio</i>	0.04	0.12	0.1	9
Polychaeta				
<i>Tomopteris helgolandica</i>	0.15	0.20	2.5	36
<i>T. elegans</i>	0.76	0.75	13.1	100
<i>Vanadis crystallina</i>	0.04	0.12	0.6	9
<i>Callizonella lepidota</i>	0.11	0.19	1.9	27

<i>Pelagobia longicirrata</i>	3.64	3.93	63.1	72
<i>Sagittella kowalevskii</i>	1.05	1.46	1.9	18
<i>Travisopsis lanceolata</i>	0.29	0.74	16.3	45
<i>Typhloscolex muelleri</i>	0.04	0.12	0.6	9
Chaetognatha				
<i>Sagitta enflata</i>	45.09	68.78	24.7	100
<i>S. minima</i>	122.11	88.72	67.1	100
<i>S. setosa</i>	1.75	1.61	1.0	81
<i>S. serratodentata</i>	4.15	6.36	2.3	100
<i>S. bipunctata</i>	0.11	0.19	3.7	27
<i>S. decipiens</i>	0.41	0.80	0.2	27
<i>S. lyra</i>	1.56	2.99	0.9	45
<i>Krobnitta subtilis</i>	0.04	0.12	0.1	9

Table 3. Pearson coefficient of correlation between abundance of dominant species and hydrographic parameters ($n = 11$). Asterisks indicate the level of significance: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Taxa	Temperature	Salinity
Species		
<i>Aglaura hemistoma</i>	-0.348	-0.243
<i>Liriope tetrphylla</i>	0.444	0.399
<i>Lensia subtilis</i>	-0.832***	0.098
<i>Muggiaea kochii</i>	0.330	-0.551*
<i>Muggiaea atlantica</i>	-0.042	-0.152
<i>Atlanta peronii</i>	0.249	-0.288
<i>Protatlanta souleyeti</i>	0.731**	0.287
<i>Limacina inflata</i>	0.076	0.599*
<i>Creseis virgule</i>	0.297	-0.689**
<i>Creseis acicula</i>	0.247	-0.656**
<i>Pelagobia longicirrata</i>	-0.569*	-0.112
<i>Tomopteris elegans</i>	0.455	-0.192
<i>Sagitta enflata</i>	0.809***	0.276
<i>Sagitta minima</i>	0.559*	-0.393

eropoda, 8 species of Polychaeta and 8 species of Chaetognatha. Ctenophores were counted at group level and were found only sporadically. Among these species, four were recorded in the Middle Adriatic for the first time: *Muggiaea atlantica* (calycophoran), *Desmopterus papilio* (pteropod), *Protatlanta souleyeti* (heteropod), *Pelagobia longicirrata* (pelagic polychaete).

The dominant species of hydromedusae *A. hemistoma* and *L. tetrphylla* (Table 2) showed generally higher abundances in winter, spring, and autumn (Fig. 4). No correlation was found between abundances of these species

and hydrographical parameters (Table 3). Among calyco-phores, *L. subtilis*, *M. kochii* and *M. atlantica* were the most abundant and frequent species (Table 2). *L. subtilis*

attained higher abundance in winter (Fig. 4) and was in significant negative correlation with temperature ($p < 0.001$, Table 3). *M. atlantica* was most abundant in spring and attained high abundance in the Middle Adriatic (Fig. 4) very soon after its first record in the South Adriatic in February 1995 (25). Maximal abundance of *M. kochii* was registered in summer (Fig. 4). Abundance of *M. kochii* was in significant negative correlation with salinity ($p < 0.05$, Table 3). In comparison with studies before 1995, relative abundance of *M. atlantica* in calyco-phoran community markedly increased, while for *M. kochii* a decrease was registered (Table 4). The most abundant and frequent pteropods were *Limacina inflata*, *Creseis virgula* and *C. acicula* (Table 2). *L. inflata* was most abundant in autumn while *C. virgula* and *C. acicula* peaked in spring-summer period (Fig. 4). The abundances of all three species were in significant correlation with salinity ($p < 0.05$; $p < 0.01$; $p < 0.01$, respectively) (Table 3). Abundance and contribution of *C. acicula* in pteropod community was much lower than in previous investigations (Table 4). On the contrary, previously rare *L. inflata* and *C. virgula* showed relative abundance of 26 and 44% in the pteropod community, respectively (Table 4). Pteropod *Desmopterus papilio* was registered for the first time in the Middle Adriatic, in September 1995, with low relative abundance of 0.1% (Table 4).

P. longicirrata and *T. elegans* were the dominant pelagic polychaetes (Table 2). *P. longicirrata* was most abundant in winter, while *T. elegans* attained high abundance in warmer part of the year (Fig. 4). Abundances of *P. longicirrata* were in negative significant correlation




Figure 4. Abundances of dominant species in their respective groups: a) Hydromedusae, b) Calyco-phorae, c) Polychaeta, d) Pteropoda, e) Chaetognatha.

Table 4. Relative abundance (%) of newly recorded and dominant species of calyco-phorans, pteropods, heteropods and polychaetes in the Middle Adriatic in total number of specimens in the respective group (CA: calyco-phorae; PT: pteropoda; HT: heteropoda; PH: polychaeta). Comparison of our data and earlier records (references are indicated in parenthesis).

Species	Relative abundance (%) in studies earlier than 1995	Relative abundance (%) in this study (annual mean)
* <i>Muggiaea atlantica</i> (CA)	/	55%
<i>Muggiaea kochii</i> (CA)	>30% (25, 36)	12%
<i>Creseis acicula</i> (PT)	>80% (36, 37)	18%
<i>Limacina inflata</i> (PT)	<5% (36)	26%
<i>Creseis virgula</i> (PT)	<5% (36)	44%
* <i>Desmopterus papilio</i> (PT)	/	<0.1%
<i>Atlanta peronii</i> (HT)	>50% (36)	62%
* <i>Protatatlanta souleyeti</i> (HT)	/	33%
* <i>Pelagobia longicirrata</i> (PH)	+ (27)	63%
<i>Tomopteris elegans</i> (PH)	>50% (36, 38)	13%

*newly recorded species; „+” present

($p < 0.05$) with temperature (Table 3). Before 1995, *T. elegans* was the most abundant species while in this study *P. longicirrata* dominated with relative abundance of 63% in the pelagic polychaete community (Table 4).

Heteropods *Atlanta peronii* and *Protatlanta souleyeti* were the most abundant in their group but they were not so frequent in the plankton. They were registered only in June and July and from September to November, respectively (Table 2). *P. souleyeti* was not recorded previously in the Middle Adriatic and its abundance was in positive correlation with the temperature ($p < 0.01$, Table 3). *A. peronii* had high relative abundance in the heteropod community of the Middle Adriatic in earlier study as well as in this study (Table 4). In general, heteropods are not numerous, with sporadic occurrence in the Adriatic Sea (11, 33).

Among chaetognaths, *Sagitta enflata* and *S. minima* were the dominant species (Table 2). Higher abundance of *S. enflata* was found in autumn while *S. minima* peaked in warmer part of the year (Fig. 4). Abundances of *S. enflata* and *S. minima* were in significant positive correlation ($p < 0.001$ and $p < 0.05$, respectively) with temperature (Table 3).

DISCUSSION

Significant changes in the zooplankton community in the South Adriatic were recorded in 1990s, and were related to drastic changes in the thermohaline circulation of the Eastern Mediterranean (11). Similarly, in 1995/1996 notable changes in the species composition and abundances of the non-crustacean zooplankton communities in the Middle Adriatic were observed, particularly within Calycophorae, Heteropoda, Pteropoda and Polychaeta. Three newly recorded species in the Adriatic Sea, a calycophoran *Muggiaea atlantica* (25), pteropod *Desmopterus papilio*, and heteropod *Protatlanta souleyeti* (26) were found in the open Middle Adriatic in 1995. The pelagic polychaete *Pelagobia longicirrata*, registered for the first time in the North Adriatic in 1967 (27), has been found after years of absence in the South Adriatic in 1993 (26), and also during this investigation in the Middle Adriatic in 1995. All these species are common members of Atlantic Ocean fauna and have been recorded in the Western Mediterranean over past decades (28, 29, 30, 31, 32). The spreading of these species from the south towards the Middle Adriatic occurred in 1995 and coincided with the intrusion of Atlantic water (MAW) into the Adriatic Sea.

Changes of oceanographic properties of the Middle Adriatic from 1991 to 1998 are evident in higher-than-usual nutrient levels, coupled with lower-than-usual temperature, salinity and dissolved oxygen (9). These changes have been attributed to the inflow of the nutrient rich MAW into the Adriatic, caused by the anticyclonic circulation in the North Ionian Gyre. While generally lower

temperatures enabled the survival of cold-temperate species such as *M. atlantica* and *P. longicirrata*, two warm-temperate species, *D. papilio* and *P. souleyeti*, were also recorded. However, the latter two species were registered in the warmer part of the year (September, 1995) which probably enabled their survival in relatively colder conditions. Similarly, annual distributions of abundance of *P. longicirrata* and *P. souleyeti* were in strong negative and positive correlation, respectively, with temperature. Apart from species composition, abundances of previously dominant species of calycophorans, pteropods, and polychaetes have also changed dramatically. As noted in the South Adriatic from 1993 to 1996 (26, 33), the progressive dominance of *Muggiaea atlantica* over the formerly dominant congener *M. kochii* was registered, the pteropod *Creseis virgula* supplanted *C. acicula* and the previously very rare *Pelagobia longicirrata* became dominant pelagic polychaete in the Middle Adriatic in 1995/1996. The analysis of long-term zooplankton data series from the Stončica station (1960–1992) indicated the onset of changes in the abundance of some gelatinous zooplankton from 1980s at station Stončica (19). In addition, from mid '80s, to mid '90s, changes were observed in the structure and dynamics of the Northern Adriatic copepod community (34) as well as in the microzooplankton community of the South Adriatic (35). According to the long-term zooplankton data (1993–2011) from the South Adriatic, *M. atlantica*, *P. longicirrata* and *P. souleyeti* became established in the Adriatic zooplankton community while the presence of *D. papilio* depended of the type of current that prevailed in the Adriatic (11).

All these faunal changes can be associated with the change of North Ionian Gyre circulation and, consequently with inflow of Atlantic Water (MAW) into the Adriatic which was uniquely strengthened by the Eastern Mediterranean Transient (EMT) in the early 1990s. The magnitude of influence of this phenomenon on the Mediterranean ecosystem is still not completely known which makes presented data essential in the evaluation of biological changes in the Adriatic Sea. Additionally, these findings will allow us to track future changes in the Adriatic pelagic community in the light of the BiOS theory and circulation changes in the NIG on decadal time scale.

REFERENCES

- OVCHINNIKOV I M, ZATS V I, KRIVOSHEYA V G, UDO-DOV A I 1985 A forming of deep eastern Mediterranean water in the Adriatic Sea. *Okeanologia* 25: 911–917 (in Russian)
- BORZELLI G L E, GAČIĆ M, CARDIN V, CIVITARESE G 2009 Eastern Mediterranean Transient and reversal of the Ionian Sea circulation. *Geophys Res Lett* 36: L15108 <http://dx.doi.org/10.1029/2009GL039261>
- GAČIĆ M, BORZELLI G L E, CIVITARESE G, CARDIN V, YARI S 2010 Can internal processes sustain reversals of the ocean upper circulation? The Ionian Sea example. *Geophys Res Letters* 37: L09608

4. GAČIĆ M, SCHROEDER K, CIVITARESE G, COSOLI S, VETRANO A, BORZELLI G L E 2013 Salinity in the Sicily Channel corroborates the role of the Adriatic–Ionian Bimodal Oscillating System (BiOS) in shaping the decadal variability of the Mediterranean overturning circulation. *Ocean Sci* 9: 83–90. <http://dx.doi.org/10.5194/os-9-83-2013>
5. CIVITARESE G, GAČIĆ M, LIPIZER M, BORZELLI G L E 2010 On the impact of the bimodal oscillating system (BIOS) on the biogeochemistry and biology of the Adriatic and Ionian Seas (Eastern Mediterranean). *Biogeosciences* 7: 3987–3997. <http://dx.doi.org/10.5194/bg-7-3987-2010>
6. ROETHER W, KLEIN B, MANCA B B, THEOCHARIS A, KIROGLOU S 2007 Transient Eastern Mediterranean deep waters in response to the massive dense-water output of the Aegean Sea in the 1990s. *Progress Oceanogr* 74: 540–571
7. KLEIN B, ROETHER W, MANCA B B, BREGANT D, BEITZEL V, KOVAČEVIĆ V, LUCHETTA A 1999 The large deep water transient in the Eastern Mediterranean. *Deep-Sea Res Pt. I*, 46: 371–30 414
8. SAYIN E, BEŞİKTEPE T 2010 Temporal evolution of the water mass properties during the Eastern Mediterranean transient (EMT) in the Aegean Sea. *J Geophys Res* 115: C10025
9. VILIBIĆ I, MATIJEVIĆ S, ŠEPIĆ J, KUŠPILIĆ G 2012 Changes in the Adriatic oceanographic properties induced by the Eastern Mediterranean Transient. *Biogeosciences* 9(6): 2085–2097. <http://dx.doi.org/10.5194/bg-9-2085-2012>
10. MIHANOVIĆ H, VILIBIĆ I, DUNIĆ N, ŠEPIĆ J 2015 Mapping of decadal middle Adriatic oceanographic variability and its relation to the BiOS regime. *J Geophys Res Oceans* 120: 5615–5630. <http://dx.doi.org/10.1002/2015JC010725>
11. BATISTIĆ M, GARIĆ R, MOLINERO J C 2014 Interannual variations in Adriatic Sea zooplankton mirror shifts in circulation regimes in the Ionian Sea. *Climate Res* 61: 231–240
12. BULJAN M, ZORE-ARMANDA M 1966 Hydrographic data on the Adriatic Sea collected in the period from 1952 through 1964. *Acta Adriat* 12: 1–438
13. BULJAN M, ZORE-ARMANDA M 1979 Hydrographic properties of the Adriatic Sea in the period from 1965–1970. *Acta Adriat* 20: 1–368
14. BULJAN M 1969 Relation between some factors affecting productivity and fish catch in the central Adriatic area. *Stud Rev Gen Fish Coun Mediterr* 41: 25–39
15. REGNER D 1981 The changes in seasonal oscillations of copepods in the central Adriatic. *Rapp Comm int Mer Médit* 27(7): 177–179
16. REGNER D 1991 Long-term investigations of copepods (zooplankton) in the coastal waters of the eastern Middle Adriatic. *Acta Adriat* 32(2): 631–740
17. BARANOVIĆ A, ŠOLIĆ M, KRSTULOVIĆ N 1993 Temporal fluctuations of zooplankton and bacteria in the middle Adriatic Sea. *Mar Ecol Prog Ser* 92: 65–75. <http://dx.doi.org/10.3354/meps092065>
18. ŠOLIĆ M, KRSTULOVIĆ N, MARASOVIĆ I, BARANOVIĆ A, PUCHER-PETKOVIĆ T, VUČETIĆ T 1997 Analysis of time series of planktonic communities in the Adriatic Sea: distinguishing between natural and man-induced changes. *Oceanologica Acta* 20 (1): 131–143
19. BERLINE L, SIOKOU-FRANGOU I, MARASOVIĆ I, VIDJAK O, FERNÁNDEZ DE PUELLES M-L, MAZZOCCHI M G, ASSIMAKOPOULOU G, ZERVOUDAKI S, FONDA UMANI S, CONVERSI A, GARCIA-COMAS C, IBANEZ F, GASPARINI S, STEMMANN L, GORSKY G 2012 Intercomparison of six Mediterranean zooplankton time series. *Prog Oceanogr* 97–100: 76–91. <http://dx.doi.org/10.1016/j.pocean.2011.11.011>
20. TISCHLER W. 1948 Biocenotische Untersuchungen an Wallhecken. *Zool. Jb* 77: 283–400
21. BODENHEIMER F. 1955 *Precis d'écologie animale*, Paris, p 315
22. BALOGH J. 1958 *Lebensgemeinschaften der Landtiere*, Berlin, p 560
23. SOUTHWOOD T R E 1975 Habitat, the templet for ecological strategies. *J Anim Ecol* 46: 337–365.
24. NAIDENOW W. 1985 Die Auswirkung der Wasserbauten auf das Zooplankton im österreichischen Donauabschnitt. – In: Naideonow W (ed) *Die Auswirkung der Wasserbaulichen Massnahmen und der Belastung auf das Plankton und das Benthos der Donau*. Verlag der bulgarischen Akademie der Wissenschaften, Sofia, 72–102
25. GAMULIN T, KRŠINIĆ F 2000 Calyctophores (Siphonophora, Calyctophora) of the Adriatic and Mediterranean Seas. *Nat Croat* 9: 1–198
26. BATISTIĆ M, KRŠINIĆ F, JASPRICA N, CARIĆ M, VILIBIĆ I, LUČIĆ D 2004 Gelatinous invertebrate zooplankton of the South Adriatic: species composition and vertical distribution. *J Plankton Res* 24: 459–474. <http://dx.doi.org/10.1093/plankt/fbh043>
27. POŽAR A 1972 Polychaeta obraštajnih životnih zajednica na različitim podlogama. *Rad JAZU* 364: 39–46 (in Croatian)
28. FURNESTIN M L 1960 Zooplankton du Golfe du Lion et de côte orientale de Corse. *Rev Trav Inst Pêches Marit* 24: 153–252
29. DI GERONIMO I 1970 Heteropoda e Pteropoda Thecosomata in sedimenti abbissali recenti dello Jonio. *Thalassia Salent* 4: 41–115
30. RAMPAL J 1975 Les Thécosomes (Mollusques pélagiques). *Systématique et évolution – Écologie et biogéographie méditerranéennes*. PhD dissertation, l'Université de Provence, Aix-Marseille, p 485 (in French)
31. PLEIJEL F, DALES R P 1991 Polychaetes: British Phyllococoidaeans, Typhloscolecoideans and Tomopteroideans. In: Kermack DM, Barnes RSK (eds) *Synopses of the British Fauna (New Series)*, 45. Universal Book Services, Oegstgeest, p 202
32. PINCA S, DALLOT S 1995 Meso- and macrozooplankton composition patterns related to hydrodynamic structures in the Ligurian Sea (Trophos-2 experiment, April–June 1986). *Mar Ecol Prog Ser* 126: 49–65. <http://dx.doi.org/10.3354/meps126049>
33. BATISTIĆ M, JASPRICA N, CARIĆ M, LUČIĆ D 2007 Annual cycle of the gelatinous invertebrate zooplankton of the eastern South Adriatic coast (NE Mediterranean). *J Plankton Res* 29: 671–686. <http://dx.doi.org/10.1093/plankt/fbm048>
34. KAMBURSKA L, FONDA-UMANI S 2006 Long-term copepod dynamics in the Gulf of Trieste (Northern Adriatic Sea): recent changes and trends. *Clim Res* 31: 195–203. <http://dx.doi.org/10.3354/cr031195>
35. KRŠINIĆ F 2010 Tintinnids (Tintinnida, Choreotrichia, Ciliata) in the Adriatic Sea, Mediterranean. Part I. Taxonomy. Institute for Fisheries and Oceanography, Split, p 186
36. GAMULIN T 1979 Zooplankton istočne obale Jadranskog mora. *Acta biol* 8:177–270 (in Croatian)
37. LUČIĆ D, MIKUŠ J 1995 Mrežni zooplankton. In: Zvonarić, T (ed.) *Kontrola kvalitete obalnog mora*. Projekt Vir-Konavle 1994. IOR, pp 67–80.
38. ZEI, M 1956 Pelagic polychaetes of the Adriatic. *Thalassia Jugosl* 1: 33–68