

Microfacies characteristics of carbonate cobble from Campanian of Slovenj Gradec (Slovenia): implications for determining the *Fleuryana adriatica* De Castro, Drobne and Gušić paleoniche and extending the biostratigraphic range in the Tethyan realm

The Mining-Geology-Petroleum Engineering Bulletin
UDC: 551.7:551.8:56
DOI: 10.17794/rgn.2018.4.1

Original scientific paper



Alan Moro¹, Ivo Velić², Vasja Mikuž³, Aleksander Horvat⁴

¹ Department of Geology, Faculty of Science, University of Zagreb, Horvatovac 102A, Zagreb, amoro@geol.pmf.hr

² Croatian Geological Survey, Sachsova 2, Zagreb; Croatian Geological Summer School, Zagreb, ivo.velic@zg.t-com.hr

³ University of Ljubljana, Department of Geology, Privoz 11, SI-1000 Ljubljana, Slovenia, vasja.mikuz@ntf.uni-lj.si

⁴ Research Centre of the Slovenian Academy of Sciences and Arts, Ivan Rakovec Institute of Palaeontology, Novi trg 2, SI-1000 Ljubljana, Slovenia, ahorvat@zrc-sazu.si

Abstract

A carbonate cobble was collected in a creek near Slovenj Gradec (Slovenia). It comprises *Trochactaeon* gastropods as well as foraminifera *Fleuryana adriatica* and *Cuneolina ketini* from a subtidal environment with agglutinated and porcelaneous benthic foraminifera. Packstone-grainstone of the structural type contains the abovementioned foraminiferal taxa, which are previously undetermined in this part of the Upper Cretaceous Tethyan realm. Stratigraphic data from this area, together with the determined microfossil assemblage of the carbonate cobble, indicate the upper part of Lower Campanian age, implying an extension of the *F. adriatica* and *C. ketini* stratigraphic range. The paleoniche within the Tethyan realm in which *F. adriatica* thrive is the shallowest part of the subtidal, close to peritidal conditions. It was developed on a relatively harder substrate that resulted from a laterally more pronounced paleorelief which enabled the shifting of hyaline benthic foraminifers towards the shallowest part of the subtidal. Their paleoniche is laterally absent from protected paleoenvironments with soft muddy substrate and a relatively high carbonate sedimentation rate, colonized with elevator rudists, as well as from relatively higher water energy paleoenvironments with completely hard grainstone-rudstone substrate.

Keywords:

Lower Campanian, *Fleuryana adriatica*, *Cuneolina ketini*, benthic foraminifera, paleoniche, carbonate deposits

1. Introduction

Near the Upper Cretaceous Slovenj Gradec locality, carbonate cobble rich in Gosau-type gastropod fauna (see Figure 1) was collected from a creek during previous field investigations (Moro et al., 2016). In a paper by Mikuž et al. (2012), typical Gosau gastropods *Trochactaeon* cf. *goldfussi* (d'Orbigny) and *Trochactaeon giganteus* (Sowerby) were determined. To determine microfacies characteristics, a single thin section was also made from the carbonate cobble with trochactaeons. Within the agglutinated-porcelaneous foraminiferal assemblage, *Fleuryana adriatica* De Castro, Drobne, and Gušić was determined, biostratigraphically important for the Upper Campanian to Late Maastrichtian (De Castro et al. 1994; Velić 2007; Fleury, 2014; Solak et al., 2017). Determination of *F. adriatica* suggests a younger age of the eroded carbonate cobble compared with the nearby Slovenj Gradec succession, which has a chronostratigraphic age of the upper part of the Lower Campanian (Moro et al., 2016). Considering the other

Gosau-type sedimentary successions where Campanian acteonelloids always lie below rudists bearing limestones (Sanders and Baron Szabo, 1997; Kollmann, 2014), the bistratigraphical range of *F. adriatica* (Upper Campanian to Late Maastrichtian) (Fleury, 2014; Solak et al. 2017) needs to be reconsidered.

Latest Jurassic to Early Cretaceous orogen convergence (Marton et al., 2017) was thrusted in the investigated area, which was overridden by a nappes system. Subsequently, large parts of the orogen were uplifted and subareally eroded (Ratschbacher et al., 1989, Sanders et al., 2004). A characteristic of the Late Cretaceous tectonics of the Alpine-Carpathian-Pannonian part of the Tethyan realm is the synchronous formation of Gosau-type sinsedimentary basins. Initial subsidence from Cenomanian to Campanian times within Gosau-type basins is associated with the development of a fluvial-lacustrine to shallow marine environment and the deposition of conglomerates, coal-bearing marls, siliciclastics rich in corals and acteonelloid gastropods and rudist limestones on the top of the succession (Willing-shofer et al., 1999). As a result of the oblique orogenic convergence, the abovementioned deposits were mainly

Corresponding author: Alan Moro
amoro@geol.pmf.hr



Figure 1. Photographs (above, below and from the side) of the carbonate cobble

controlled by strike-slip extensional tectonics (**Willinghofer et al., 1999; Faupl and Wagreich, 2000; Sanders et al., 2004**). Emersion, regression, and Miocene transgression (**Mioč, 1978; Brlek et al. 2016; Sremac et al., 2016; 2018; Bošnjak et al., 2017**) hide Upper Cretaceous outcrops and make them hard to find for possible paleoreconstruction.

This paper aims to resolve the question of the stratigraphical disparity between the carbonate cobble and the Slovenj Gradec succession, as well as to determine the extension of the biostratigraphical range and paleoecological niche of the *Fleuryana adriatica* microfossil association for the Tethyan realm.

2. Geological background of *Fleuryana adriatica* and the associated microfossil assemblage

Upper Cretaceous Tethyan localities with findings of *F. adriatica* and associated microfossils are rare, and research studies range from the description of a new species and determination of associated microfossils to paleoenvironmental reconstructions and chronostratigraphical determinations.

In the Maastrichtian of the Latium area (Italy) (**Chiocchini and Mancinelli, 2001**), *F. adriatica* occurs within a vertical exchange of laminated dolomites and partly dolomitized mudstones/wackestones of “Discorbidae e Miliolidae biozone” (**Chiocchini and Mancinelli, 2001; Chiocchini et al., 2012**). Associated microfossils are mostly composed of Ostracoda, Miliolidae, and Charophyta, while *Laffitteina marsicana*

Farinacci, *Fleuryana adriatica* De Castro Drobne and Gušić and *Rhapydionina liburnica* Stache are present only in the upper part of the unit. The biostratigraphic age is determined as the uppermost Maastrichtian.

In Central Anatolia of Turkey, the paleoenvironment of *C. ketini* is described as reefal limestones with basal transgressive conglomerates, laterally passing into ophiolitic conglomerate, pelagic limestones, and limestones with chert nodules (**Inan, 1988**). The associated microfossil assemblage consists of *Loftusia minor* Cox, *L. harrisoni* Cox, *Siderolites calcitrapoides* Lamarck, *Omphalocyclus macroporus* (Lamarck), *Orbitoides media* (D’Archiac), *O. apiculatus* Schlumberger, *Torreina torrei* Palmer, *Pseudomphalocyclus blumenthalii* Meriç, *Cuneolina* sp. and *Operculina* sp. According to the associated hyaline larger benthic foraminifers, the chronostratigraphic range of *C. ketini* is determined as the Upper Maastrichtian (**Inan, 1988**).

In Haftoman Formation of Central Iran (**Wilmsen et al., 2018**), representing an eperic carbonate platform, *C. ketini* thrives in a low- to moderate-energy paleoenvironment of an outer platform characterized by bioclastic rudstone/floatstone/boundstone. Associated fossils are *Nezzazatinella* sp., textularids, miliolids, coiled recumbent rudists, echinoderms, and ostracods green algae (dasy-clad/*Cymopolia*? sp.). This paleoenvironment is in a lateral exchange with (1) baffles with small elevator rudists and red algae, (2) bioclastic rudstones with orbitoids, miliolids, inoceramids, rudists fragments, udotea-can green algae, red algae, and gastropods, and (3) bindstone with red algae, fragments of corals, bryozoans, rudists, thick-shelled bivalves, *Orbitoides media* (D’Archiac), *Praesiderolites douvillei* Wannier, and miliolid foraminifera. The presence of hyaline larger benthic foraminifers suggests a biostratigraphic range of *C. ketini* as Middle to Late Campanian (**Wilmsen et al., 2018**).

In western Central Taurides, the paleoenvironment of *F. adriatica* is peritidal of a restricted platform without any siliciclastic input and the occasional influence of more open water circulation (**Solak et al., 2017**). The associated microfossil assemblage consists of *Moncharmontia apenninica* (De Castro), *Murgeina* sp., *Arenobulimina* sp., *Bolivinopsis* sp., *Rotalispira scarsellai* (Torre), *Nezzazatinella* sp., *Nezzazata* spp., *Minouxia* sp., *Stensioeina surrentina* Torre, *Accordiella conica* Farinacci, *Scandonea samnitica* De Castro, *Murciella* gr. *cuvillieri* Fourcade, *Biconcava* sp., *Pseudocyklamina sphaeroidea* Gendrot, *Rhapydionininae*, *Cuneolina* gr. *pavonia* D’Orbigny, discorbids, miliolids, Nubeculariidae, algae (*Thaumatoporella parvovesiculifera* Raineri, Dasycladaceae and charophyta oogonia), cyanobacteria *Gahkumella huberi* Zaninetti, *Decastronema kotori* Radoičić, and ostracods. Associated macrofossils are rudists, mainly radiolitids and gastropods. The biostratigraphic range is from the Upper Campanian (smaller-size specimens) to the Maastrichtian (identical to typical specimens) (**Solak et al., 2017**).



Figure 2. Microphotographs of microfacies. a.- Laminated and partly bioturbated packstone-grainstone; b. - Packstone-grainstone with rare fragments of macrofossils; c, d. - Floatstone rudstone with fragments of macrofossils, abundant miliolids and *Dicyclina* (d); e, f. - Small gastropod within floatstone – rudstone with miliolids and *Nezazatinella* (e)

In the Zagros zone in SW Iran (**Schlagintweit and Rashidi, 2016**), *F. adriatica* thrives in foraminifer-dasycladacean soft substrate wackestones-packstones and the very low water energy of the shallow protected inner platform area (lagoon). The associated microfossil assemblage consists of *Omphalocyclus macroporus* (Lamarck), *Spirolina farsiana* Schlagintweit and Rashidi, *Loftusia* spp., *Accordiella tarburensis* Schlagintweit and Rashidi, *Dicyclina schlumbergeri* Munier-Chalmas, *Broeckinella arabica* Schlagintweit and Rashidi, *Gyro-*

conulina columellifera Schroeder & Darmoian, *Neobalkhania bignoti* Cherchi, Radoičić, & Schroeder, and *Laffiteina monodi* Marie. Dasycladales are represented by *Pseudocymoplia anadyomenea* (Elliott), *Pseudocymoplia?* sp., and *Salpingoporella pasmanica* Radoičić. Rudists are present through broken shells within packstone/grainstone-floatstone/rudstones. Besides the abovementioned taxa, in the shallowest, upper part of the succession, the foraminiferal association comprises: *Antalyina korayi* Farinacci & Köylüoglu, *Cuneolina* sp.,




Figure 3. Microphotographs of *C. ketini* and associated benthic foraminifers. a. - *Accordiella conica* Farinacci; b., h., - *Dicyclina schlumbergeri* Munier-Chalmas and *Nezzazatinella* sp. (b); c. - *Cuneolina parva* Henson; d, k- *Cuneolina* sp.; i, j. - *Cuneolina ketini* Inan; e - *Moncharmontia compressa* (De Castro); f, g. - *Moncharmontia appenninica* (De Castro)

Dictyoconella minima Henson, *Elazigella?* sp., *Fallotia* aff. *jacquoti* Douvillé, *Loftusia harrisoni* Cox, *Loftusia coxi* Henson, *Loftusia morgani* Douvillé, *Mississippina?* *binkhorsti* (Reuss), *Nezzazatinella?* cf. *picardi* (Henson), *Neobalkhania bignoti* Cherchi, Radoičić & Schroeder, *Elphidiella?* *cretacea* (Pérébaskine), Orbitoliniidae indet., *Tarburina zagrosiana* Schlagintweit, Rashidi & Barani, and *Valvulina?* sp. The biostratigraphic age is determined as the Maastrichtian (**Schlagintweit and Rashidi, 2016**).

At localities from the Gavrovo-Tripolitza zone (Greece) (**Fleury, 2014**), *F. adriatica* thrive within an Upper Campanian to Late Maastrichtian inner platform paleoenvironment characterized by the vertical exchange of subtidal and peritidal conditions together with *Cuvillierina salentina* Papetti & Tedeschi, *Murciella cuvillieri* Fourcade, *Murciella renzi* Fleury, *Cyclopseudedomia smouti* Fleury, *C. hellenica* Fleury, *C. klokovaensis* Fleury, and *Rhapydionina dercourtii* Fleury. In rare places, this paleoenvironment is in a lateral exchange with deeper, more open subtidal conditions with hyaline benthic foraminifers (**Fleury, 2014**).

In the Gosau-type sedimentary succession, *C. ketini* and *F. adriatica* were first mentioned by **Sanders et al. (2004)** from the Campanian deposits of the Central Alps.

The paleoenvironment of *C. ketini* and *F. adriatica* is determined as an external part of the platform. Associated microfauna comprise *Accordiella conica* Farinacci, *Bolivinopsis* sp., *Calveziconus* cf. *leacalvezae* Caus & Cornellà, *Cuneolina* sp., *Dicyclina schlumbergeri* Munier-Chalmas, *Fleuryana* sp., *Lenticulina* sp., *Marssonella oxycona* (Reuss), *Minouxia* sp., *Mississippina* sp., *Orbitoides* sp., *Pseudosiderolites* sp., and *Reticulinella fleuryi* Cvetko, Gušić & Schroeder. Associated macrofossils are Hippuritidae (*Vaccinites vesiculosus* (Woodward), *V. ultimus* (Milovanović), *Hippurites colliciatus* Woodward, Radiolitidae (*Radiolites* cf. *angeioides* (Lapeirouse), *Pseudopolyconites* sp. and *Joufia* sp.) as well as fragments of corals, echinoids, coralline algae, brachiopods, non-rudist bivalves, bryozoans, skeletal sponges, and calcareous green algae. The chronostratigraphic range is determined as Late Campanian (**Sanders et al., 2004**).

At Maastrichtian deposits on Brač Island, the paleoenvironment of *F. adriatica* (**De Castro et al., 1994**; **Korbar et al., 2017**) is subtidal with *Bolivinopsis* sp., *Laffitteina mengaudi* (Astre), primitive “agglutinated” conical foraminifera, abundant miliolids, nezzazatids, Discorbidae, Rotaliidae, Ophthalmidiidae, Valvulinidae, associated with Ostracoda, remains of algae *Thaumato-porella parvovesiculifera* (Raineri) and charophytes, requieniid rudists *Apricardia* sp., and rare planktonic fo-




Figure 4. Microphotographs of associated benthic foraminifers. a, e. - *Dicyclina schlumbergeri* Munier-Chalmas; b, d, h. - *Dictyopselloides cuvillieri* (Gendrot); c. - *Cuneolina parva* Henson; f, g, l. - *Pseudocyclammina sphaeroidea* Gendrot and *F. adriatica* (g); i, j, k. - *Dictyopsella kiliani* Munier-Chalmas

raminifera and pithonellid calcispheres (Korbar et al., 2017). The chronostratigraphic range is determined by comparison with the type-locality (De Castro et al., 1994), as well as the Sr isotope stratigraphy data of Steuber et al. (2005), which confirm the terminal Maastrichtian age.

Within the carbonate facies of the Slovenj Gradec succession, characterized by packstone-grainstone and floatstone-rudstone textural types, completely preserved macrofossils are rare and consist of small radiolariid individuals in the succession's lower part. Fossil assemblage comprises the following agglutinated and porcellaneous foraminiferal species: *Accordiella conica* Farinacci, *Calveziconus* cf. *lecalvezae* Caus & Cornella, *Scandinea samnitica* De Castro, *Dicyclina schlumbergeri* Munier-Chalmas, *Dictyopsella kiliani* Munier-Chalmas, *Cuneolina pavonia* D'orbigny and *Pseudocyclammina sphaeroidea* Gendrot (Moro et al., 2016). Determined benthic foraminifers indicate the *Calveziconus* cf. *lecalvezae* subzone (Frijia et al., 2015), which chronostratigraphically corresponds to the upper part of the Lower Campanian (Moro et al., 2016). According to the benthic fossil assemblage and textures, the depositional environment can be interpreted as shallow subtidal with moderate water energy.

3. Methods

Thin-section analyses of the limestone cobble as well as the Slovenj Gradec succession include textures and skeletal components and biostratigraphic characteristics. Visual percentage charts were used to estimate the relative abundance of grains (Bacelle and Bosellini, 1965; in Flügel, 2004), and emphasize any differences in microfacies of the carbonate cobble as well as microfacies differences between the cobble and the Slovenj Gradec succession. The taxonomic study of benthic foraminifers is based on randomly orientated sections of tests with observable morphological characteristics of different species. The taxonomic framework used to identify species builds on papers by Inan (1988), De Castro et al. (1994), and Velić (2007).

4. Results

4.1. Microfacies characteristics and chronostratigraphy of the carbonate cobble

The studied cobble is entirely composed of carbonate, with dimensions 20x10 cm. At the macroscopic level, it shows a uniform texture without any visible structure.




Figure 5. *Fleuryana adriatica*. Axial (e, f, g, h, i), equatorial (a, b, d, k, l), and oblique (c, j, m, n, o, p) sections

Macroscopic two cross-sections of Gosau-type gastropods are visible (see **Figure 1**), determined as *Trochactaeon giganteus* (Sowerby) and *Trochactaeon* cf. *goldfussi* (d'Orbigny). A detailed description of the mollusk fauna is given in **Mikuž et al. (2012)**. Thin sections, 56 in total, reveal shallow-water non-skeletal and skeletal particles, including pellets, peloids, benthic foraminifers, and fragments of macrofossils, presented through two textural types: packstone-grainstone and floatstone-rudstone (see **Figure 2**), which appears randomly throughout the thin sections. Packstone-grainstone characterizes bioclasts (10-20%) with similar frequencies of shallow water macrofossil fragments (5-10%) and whole microfossils (5-10%). Within the microfossil assemblage, miliolids (2.5-5%) predominate over cuneolinids and rare dicyclinas (1 to 2.5%). The frequency of small nezzazatids (*Fleuryana* and *Moncharmontia*: **Loeblich and Tappan, 1988; Kaminski, 2004**, included within the family Charentiidae) ranges up to 1%. The frequency of *Nezazatinella* is also up to 1%, as well as test sections of undetermined benthic forams (see **Figure 4I**) with a frequency of up to 1%. Within estimated frequency ranges, some foraminifera species are present rarely:

Accordiella conica and *Moncharmontia compressa* represented by a single specimen, *Pseudocyklamina sphaeroidea* and *Cuneolina ketini* by two specimens, and *Moncharmontia appeninica* by three specimens.

The microfossil assemblage is composed of the following agglutinated and porcellaneous foraminiferal species (see **Figures 3 to 7**): *Accordiella conica* Farinacci, *Cuneolina ketini* Inan, *Fleuryana adriatica* De Castro, Drobne & Gušić, *Dicyclina schlumbergeri* Munier-Chalmas, *Dictyopsella kiliani* Munier-Chalmas, *Dictyopselloides civilieri* (Gendrot), *Cuneolina parva* Henson, *Moncharmontia appeninica* (De Castro), *M. compressa* (De Castro) and *Pseudocyklamina sphaeroidea* Gendrot.

The determined benthic foraminifers as well as the microfossil assemblage of the nearby Slovenj Gradec section indicate the *Calveziconus* cf. *lecalvezae* subzone (**Frijia et al., 2015**), which chronostratigraphically correspond to the upper part of the Lower Campanian (**Moro et al., 2016**) (see **Figure 8**). According to the benthic fossil assemblage, structures, and textures, the depositional environment can be interpreted as shallow subtidal with moderate water energy.




Figure 6. *Fleuryana adriatica*. Axial (n), equatorial (d, i, k) and oblique (a, b, c, d, e, f, g, h, j, l, m, o, p) sections

4.2. Microfacies characteristics of the Slovenj Gradec succession

The carbonate facies of the Slovenj Gradec succession are packstone-grainstone and floatstone-rudstone, with both lithotypes distributed randomly throughout the succession. Completely preserved macrofossils are rare and consist of small radiolarid individuals up to 4 cm in height and 0.8 cm in diameter in the succession's lower part. Thin sections, 28 in total, reveal shallow-water non-skeletal and skeletal particles, including peloids, benthic foraminifers, and fragments of macrofossils. Within the packstone-grainstone, the estimated frequency of bioclasts is 2.5 to 20%, where fragments of macrofossils are present with a frequency of 5 to 20%, while microfossils are present with a frequency of up to 2.5%, with a random domination of miliolids or cuneolinids and rare dicyclinids. Other microfossils (such as small nezazatinellids, *Accordiella conica*, *Calveziconus* cf. *lecalvezae*, *Dictyopsella kiliani* and previously undetermined *Nezzazatinella* sp.) are present in small numbers, from 1 to 4 specimens. Within the floatstone-rudstone variates, the estimated frequency of bioclasts, rarely

lithoclasts, is between 50 and 60%. Fragments of radiolitids predominate while microfossils are present with a small number of miliolids and nezazzatids.

5. Paleoniche of *Fleuryana adriatica*

The subtidal carbonate paleoenvironment with "typical" biostratigraphically important, mostly epiphytic (**Murray, 2006**), porcelaneous-agglutinated-hyaline foraminifers (such as *Scandonea*, *Moncharmontia*, *Accordiella*, *Calveziconus*, *Dicyclina*, *Dictyopsella*, *Orbitoides*, *Praesiderolites*) and rudists, appears at the Tethyan realm during the Campanian-Maastrichtian in textural types, ranging mostly from floatstone-rudstone to wackestone-packstone (**Gušić and Jelaska, 1990**; **Moro et al., 2002, 2016**; **Chiocchini and Mancinelli, 2001**; **Steuber et al., 2005**; **Velić, 2007**; **Schlagintweit and Rashidi, 2016**; **Solak et al., 2017**). Although there are 10 cycle boundaries, i.e. beginnings and terminations of sequences towards the end of the Cretaceous (see **Figure 8**), which could allow the possibility of the development of shallow-water paleoenvironments with the above-mentioned benthic foraminifers, probably a tectonic




Figure 7. *Fleuryana adriatica*. Axial (b, j), equatorial (a, c, d, e, g, i, l, m, r, t) and oblique (f, h, k, n, o, p, s) sections

overprint of the sea-level change resulted in the regional development of paleogeographically and biostratigraphically more or less extended subtidal paleoenvironments during the Campanian and Maastrichtian (Gušić and Jelaska, 1990; De Castro et al., 1994; Chiocchini and Mancinelli, 2001; Moro et al., 2002, 2016; Vlahović et al., 2005; Steuber et al., 2005; Steuber and Schüter, 2012; Schlagintweit and Rashidi, 2016; Korbar et al., 2017; Solak et al., 2017). Somewhere within that paleoenvironment is paleoniche where *F. adriatica* thrived.

Similarly to the distribution of benthic foraminifers (Hottinger, 1983), elevator rudist morphotypes thrive in soft muddy substrate environments, which range from the shallowest subtidal with a pure radiolarid assemblage over a relatively deeper (absence of shallowing upward cycles) subtidal with a mixed radiolarid-hippuritid assemblage towards a subtidal with relatively rare vaccinitid individuals (Gušić and Jelaska, 1990; Moro, 1997; Moro and Čosović, 2000; Moro et al., 2002, 2016; Steuber et al., 2005; Korbar et al., 2009). Exceptions are radiolarids, which could also thrive in shallow-water, moderate-energy paleoenvironments as clingers (Ross and Skelton, 1993; Skelton and Gili, 2002).

The rudist assemblage, which thrives in a subtidal paleoenvironment as elevators or occasionally as clingers (radiolarids) (Skelton and Gili, 2002) with mostly epiphytic porcelaneous-agglutinated foraminifers (Murray, 2006), could also have laterally overlapped with hyaline foraminifers (Gušić and Jelaska, 1990; Sanders et al., 2004; Schlagintweit and Rashidi, 2016; Solak et al., 2017; Wilmsen et al., 2018) on a harder substrate (Hottinger, 1983).

The presence or absence of *F. adriatica* and associated microfossils within that kind of shallow subtidal environment (Inan, 1988; De Castro et al., 1994; Chiocchini and Mancinelli, 2001; Sanders et al., 2004; Velić, 2007; Fleury, 2014; Schlagintweit and Rashidi, 2016; Solak et al., 2017; Korbar et al., 2017; Wilmsen et al., 2018) (see Figure 9) is presumably the result of the lateral presence of a relatively harder substrate (Hottinger, 1983), more adequate for epiphytic benthic foraminifers (Murray, 2006). This lateral variation could be the result of slightly pronounced uplifts or relatively steeper parts of the paleorelief within a more open subtidal. This kind of substrate (sea floor) even includes peritidal paleoenvironments with shallowing upward cy-




Figure 8. Biostratigraphic and chronostratigraphic ranges of microfossils. Legend: 1 – Biostratigraphic ranges after **Velíć** (2007), **Solak et al.** (2017) and **Wilmsen et al.** (2018). 2 – Chronostratigraphic ranges after **Steuber et al.** (2005), **Frijia et al.** (2015) and **Moro et al.** (2016). 3 - Extended biostratigraphic ranges of *F. adriatica* and *C. ketini* (this paper). 4 – Chronostratigraphically extended ranges on genera level after **Moro et al.** (2016). The horizontal line indicates the beginning of the transgression of the Slovenj Gradec succession (after **Moro et al.** 2016). Sea-level curve and sequences after **Haq** (2014). Rudist biozones after **Steuber and Schlüter** (2012). Microfossils determined in the carbonate cobble and the S. Gradec succession are marked with a star.

cles and a sea floor hard enough for the presence of hyaline benthos (**Schlagintweit and Rashidi, 2016; Solak et al., 2017**). The moderate hydrodynamically energized packstone-grainstone texture enables the shifting of epiphytic (**Murray, 2006**) hyaline foraminifers towards shallower parts of the subtidal (**Hottinger, 1983; Hohenegger, 1999; Hohenegger, 2000**) (see **Figure 9**) where they overlap with the porcelaneous-agglutinated benthic assemblage (**Velić, 2007**). Similarly, sediment dwelling rudists are pushed towards the mud-prevailing shallower parts of the subtidal (**Sanders et al., 2004; Schlagintweit and Rashidi, 2016; Solak et al., 2017; Wilmsen et al., 2018**). Moreover, *F. adriatica* paleo-niche is laterally absent from protected paleoenvironments with soft muddy substrate and a relatively high carbonate sedimentation rate, colonized with elevator radiolitids and hippuritids (**Gušić and Jelaska, 1990; Chiocchini and Mancinelli, 2001; Wilmsen et al., 2018**) as well as from the relatively higher water energy paleoenvironments characterized by hard grainstone-

rudstone substrate (Gušić and Jelaska, 1990; Schlüter et al., 2008; Wilmsen et al., 2018).

Besides the type of substrate, salinity, turbidity, and nutrient supply could also be controlling factors for the macro and microfauna of the Upper Cretaceous Tethyan realm with subtropical conditions (Camoin et al., 1993). The abundant presence of miliolids (De Castro et al., 1994; Chiocchini and Mancinelli, 2001; Solak et al., 2017; Korbar et al., 2017) and gastropods, typical of shallow-water environments, could also indicate elevated (Murray, 2006) meso to eurihaline conditions (Kollmann, 1965; Herm, 1977; Sohl and Kollmann, 1985). The lateral presence of brackish to fresh-water Characean algae (Korbar et al., 2017) suggests at least the occasional influence of fresh water, while the rare planktonic foraminifera and calcispheres indicate a more open depositional environment (Korbar et al., 2017; Solak et al., 2017). Their presence could exclude salinity as a limiting factor. Turbidity is less likely a controlling factor within moderate-energy shallow-water environments




Figure 9. Block diagram with reconstructed paleoniche of *Fleuryana adriatica*. Not to scale. Explanation: 1-3 benthic foraminifers: 1.- hyaline, 2.- porcelaneous, 3.- agglutinated, 4.- planktonic foraminifers, 5-9 depositional environments: 5.- peritidal, 6.- shallow subtidal, 7.- deeper subtidal, 8.- deepwater (pelagic), 9.- deepwater marly sediments, 10.- paleoniche of *F. adriatica*.

where gastropods, rudists, and corals once thrived (Sanders and Baron Szabo, 1997; Moro et al., 2016). In shallow-water environments of the Tethyan realm, rudists predominate as carbonate producers, generally indicating oligotrophic conditions (Gili et al., 1995). Exceptions are successions with the presence of red algae (Sanders et al., 2004; Wilmsen et al., 2018) which may indicate, together with the large benthic foraminifers, a moderate nutrient supply (Acker and Leptoukh, 2007; Coletti et al., 2017). Although the rare local siliciclastic intercalations could reinforce nutrition as a controlling factor, as well as fragments of bryozoans (more eutrophic), corals and rudists as predominantly oligotrophic carbonate producers (Gili et al., 1995), together with the microfossil assemblage (Murray, 2006), could exclude nutrition as a controlling factor.

The presence of *F. adriatica* and other biostratigraphically important benthic foraminifers (see Figure 8) gave them the opportunity to be incorporated into a chronostratigraphic framework of rudist biozones and used in cases where fossil remnants of rudists are laterally absent.

6. Conclusions

The following conclusions may be drawn based on the microfacies analysis of the cobble from Slovenj Gradec, as well as the correlation with other localities comprising *F. adriatica* and associated microfossils within the Tethyan realm.

1. The paleoniche in which *F. adriatica* thrive is the shallowest part of the subtidal, close to peritidal conditions. The relatively harder substrate is a result of the

laterally more pronounced paleorelief. It is rare and narrow, resulting from the shifting of hard grainstone-rudstone substrate with hyaline benthic foraminifers towards the shallowest part of the subtidal. Moreover, it is laterally absent from protected paleoenvironments with soft muddy substrate and relatively high carbonate sedimentation rate, as well as from relatively higher water energy, more open paleoenvironments with hard grainstone-rudstone substrate.

2. According to the benthic microfossils and determined chronostratigraphic range of nearby successions (obtained by previous investigations), the regional biostratigraphic range of *F. adriatica* and *C. ketini* could be extended to the upper part of the Lower Campanian (80.5 Ma). Incorporated in rudist biozones and together with the ranges of other stratigraphically important foraminifers, these extended ranges could be useful in cases where fossil remnants of rudists are laterally absent.

Acknowledgments

The authors would like to thank Felix Schlagintweit and Kemal Tasli for providing valuable advice, suggestions and exact notations which improved the manuscript, as well as Jasenka Sremac and Ervin Mrnjek for helpful discussion and suggestions. The authors would also like to thank Robert Koščal for preparing the figures. The laboratory part of the research was supported by the University of Zagreb Research Programme “Sedimentology and paleontology of sedimentary successions from Outer and Inner Dinarides” while the field work was partly funded by Research Program P1-0008 financially supported by the Slovenian Research Agency.

cy. Finally, we are grateful to Murray Bales for the English proofreading.

7. References

- Acker, J.G. and Leptoukh, G. (2007): Online analysis enhances use of NASA Earth science data. Eos Transactions American Geophysical Union. 88, 14–17.
- Baccelle, L. and Bosellini, A. (1965): Diagrammi per la stima visiva della composizione precentuale nelle rocce sedimentarie. Annali della Universita di Ferrara, Sezione IX, Science Geologiche e Paleontologiche, 1, 59–62.
- Bošnjak, M., Sremac, J., Vrsaljko, D., Aščić, Š. and Bosak, L. (2017): The Miocene “Pteropod event” in the SW part of the Central Paratethys (Medvednica Mt., northern Croatia). *Geologica Carpathica*. 68, 4; 329–349. doi: 10.1515/geoca-2017-0023.
- Brlek, M., Špišić, M., Brčić, V., Mišur, I., Kurečić, T., Miknić, M., Avanić, R., Vrsaljko, D. and Slovenec, D., (2016). Mid-Miocene (Badenian) transgression on Mesozoic basement rocks in the Mt. Medvednica area of northern Croatia. *Facies*. 62, 3; doi: 10.1007/s10347-016-0470-z.
- Camoin, G., Bellion, Y., Dercourt, J., Guiraud, R., Lucas, J., Poisson, A., Ricou, L.E. and Vrielynck, B. (1993) Late Maastrichtian (69.5–65 Ma). In: Dercourt, J., Ricou, L.E., Vrielynck, B. (eds) *Atlas Tethys Palaeoenvironmental Maps, explanatory Notes*. Gauthier-Villars, Paris, pp 179–196.
- Chiocchini, M. and Mancinelli, A. (2001): *Sivasella monolateralis* Sirel and Gunduz, 1978 (Foraminiferida) in the Maastrichtian of Latium (Italy). *Revue de Micropaleontologie*, 44, 267–277.
- Chioccini, M., Pampaloni, M.L. and Pichezzi, R.M. (2012): Microfacies e microfossili delle successioni carbonatiche mesozoiche del Lazio e dell’Abruzzo Italia centrale). Memorie per servire alla descrizione della Carta Geologica d’Italia, Roma, 1-269, 156 pl.
- Coletti, G., Kateb, A.E., Bassi, D., Cavallo, A. and Spezzeri, S. (2017): Nutrient influence on fossil carbonate facies: Evidence from SEDEX extractions on Budrigalian limestones (Miocene, NW Italy and S France). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 475, 80–92. doi: 10.1016/j.palaeo.2017.03.005.
- De Castro, P., Drobne, K. and Gušić, I. (1994): *Fleuryana adriatica* n. gen., n. sp. (Foraminiferida) from the Uppermost Maastrichtian of the Brač Island (Croatia) and some other localities on the Adriatic carbonate platform. *Rasprave IV. Razreda SAZU*, 35, 129–149.
- Faupl, P. and Wagreich, M. (2000): Late Jurassic to Eocene paleogeography and geodynamic evolution of the Eastern Alps. *Mitteilungen der Österreichische Mineralogische Gesellschaft*, 92, 79–94.
- Frijia, G., Parente, M., Di Lucia, M. and Mutti, M. (2015): Carbon and strontium isotope stratigraphy of the Upper Cretaceous (Cenomanian-Campanian) shallow-water carbonates of southern Italy: Chronostratigraphic calibration of larger foraminifera biostratigraphy. *Cretaceous Research*, 53, 110–139. doi: 10.1016/j.cretres.2014.11.002
- Fleury J.-J. (2014): Données nouvelles sur *Rhaphydionina* Stache, 1913 et *Fanrhaphydionina* n. gen., un groupe de Rhaphydioninidae (Alveolinacea, Foraminifera) foisonnant en région périadiatique au Campanien-Maastrichtien. *Geodiversitas*, 36,,2, 173–208. doi: 10.5252/g2014n2a1
- Flügel, E. (2004): *Microfacies of Carbonate rocks – Analysis, Interpretation and Application*. Springer, Berlin. 976 p
- Gili, E., Masse, J.P. and Skelton, P.W. (1995): Rudists as gregarious sediment dwellers, not reef-builders, on Cretaceous carbonate platforms. *Palaeogeography, palaeoclimatology, palaeoecology*. 118, 245–267. doi: 10.1016/0031-0182(95)00064-X
- Gušić, I. and Jelaska, V. (1990): Stratigrafija gornjokrednih naslaga otoka Brača u okviru geodinamske evolucije Jadranske karbonatne platforme (*Upper Cretaceous stratigraphy of the Island of Brač within the geodynamic evolution of the Adriatic carbonate platform*). Djela Jugoslavenske akademije znanosti i umjetnosti 69, JAZU-IGI, Zagreb, 160. (in Croatian and English).
- Haq, B.U. (2014): Cretaceous eustasy revisited. *Global and Planetary change*. 113, 44–58. doi: 10.1016/j.gloplacha.2013.12.007
- Herm, D. 1977: Zyklische Regressions-Sedimentation und Fossil-Vergesellschaftungen in der Gosau (Santonium) von Brandenberg/Tirol. - Bayerische Staatssammlung für Paläontologie und historische Geologie Mitteilungen. 17, 257–277.
- Hohenegger, J. (1999): Larger Foraminifera – Microscopical greenhouses indicating shallow-water tropical and subtropical environments in the present and past. In: Hatta A, Oki K (eds) *Foraminifera as indicators of marine environments in the present and past*. Kagoshima Research Center for the Pacific Islands, Occasional Papers, 32, pp 19–45.
- Hohenegger, J. (2000): Coenoclines of larger foraminifera. *Micropaleontology*, 46, sup. 1, 127–151.
- Hottinger, L. (1983): Process determining the distribution of larger foraminifera in space and time. *Utrecht Micropaleontological Bulletin*, 30, 239–253.
- Inan, N. (1988): *Cuneolina ketini* (Foraminifere) n. sp. du Maastrichtien d’Anatolie Centrale (Turquie). *Revue de Paléobiologie*, 7, 2, 327–333.
- Kaminski, M.A. (2004): The Year 2000 classification of agglutinated foraminifera. In: Bubík, M., & Kaminski, M.A., (eds), *Proceedings of the Sixth International Workshop on Agglutinated Foraminifera. Grzybowski Revised Stratigraphic Ranges and Phanerozoic Diversity 83 Foundation Special Publication*, 8, 237–255.
- Kollmann, H. A. 1965: Actaeonellen (Gastropoda) aus der ostalpinen Oberkreide. – *Annalen des Naturhistorischen Museums in Wien* 68, 243–262, pls. 1–4.
- Kollmann, H. A. 2014: The extinct Nerineoidea and Acteonelloidea (Heterobranchia, Gastropoda): a palaeobiological approach. *Geodiversitas*, 36, 3, 349–383. doi:10.5252/g2014n3a2
- Korbar, T., Cvetko Tešović, B., Radovanović, I., Krizmanić, K., Steuber, T. and Skelton, P.W. (2009): Campanian *Pseudosabina* from the Pučišća Formation on the island of Hvar (Adriatic Sea, Croatia). *Turkish Journal of Earth Sciences*. 19, 6; 721–731. doi: 10.3906/yer-0901-9.
- Korbar, T., McDonald, I., Premec Fućek, V., Fućek, L. and Posilović, H. (2017): Post-impact event bed (tsunamite) at

- the Cretaceous–Paleogene boundary deposited on a distal carbonate platform interior. *Terra Nova*, 29, 2; 135–143 3. doi: 10.1111/ter.12257
- Loeblich, A.R. and Tappan, H. (1988): Foraminiferal genera and their classification. Van Nostrand Reinhold, New York, 970 p.
- Marton, E., Zampieri, D., Čosović, V., Moro, A. and Drobne, K. (2017): Apparent polar wander path for Adria extended by new Jurassic paleomagnetic results from its stable core: Tectonic implications. *Tectonophysics*. 700/701, 1–18. doi: 10.1016/j.tecto.2017.02.004
- Mikuž, V., Pejovnik, D., Čosović, V., Moro, A. and Horvat, A. (2012): Zgornjekredni polži Gosavskega faciesa iz okolice Slovenjega Gradca. *Folia biologica et geologica (ex Razprave IV. razreda SAZU)*. 53, 3; 5–21.
- Mioč, P. (1978): Osnovna geološka karta SFRJ 1:100.000. Tolmač za list Slovenj Gradec. Zvezni geološki zavod Beograd, 1–74.
- Moro, A. (1997): Stratigraphy and paleoenvironments of rudist biostromes in the Upper Cretaceous (Turonian-upper Santonian) limestones of southern Istria, Croatia. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 131, 113–131. doi: 10.1016/S0031-0182(96)00144-7
- Moro, A. and Čosović, V. (2000): The rudists of southern Istria – An example of environmentally induced succession within Santonian limestones. *Rivista Italiana di Paleontologia e Stratigrafia*. 106, 1, 59–71.
- Moro, A., Skelton, P.W. and Čosović, V. (2002): Palaeoenvironmental setting of rudists in the Upper Cretaceous (Turonian-Maastrichtian) Adriatic carbonate Platform (Croatia), based on sequence stratigraphy. *Cretaceous Research*. 23, 489–508. doi: 10.1006/cres.2002.1017
- Moro, A., Horvat, A., Tomić, V., Sremac, J. and Bermanec, V. (2016): Facies development and paleoecology of rudists and corals: an example of Campanian transgressive sediments from northern Croatia, northeastern Slovenia, and northwestern Bosnia. *Facies*. 62, 3; doi: 10.1007/s10347-016-0471-y.
- Murray, J. (2006): Ecology and Applications of Benthic Foraminifera, Cambridge University Press, Cambridge, 426 p.
- Ratschbacher, L., Frisch, W., Neubauer, F., Schmid, S.M. and Neugebauer, J. (1989): Extension in compressional orogenic belts: The eastern Alps. *Geology*. 17, 404–407. doi: 10.1130/0091-7613.
- Ross, D.J. and Skelton, P.W. (1993): Rudist formations of the Cretaceous: a palaeoecological, sedimentological and stratigraphical review. In: Wright, V. P. (ed.): *Sedimentology review*, 1, 73–91.
- Sanders, D. and Baron Szabo R.C. (1997) Coral-rudist bioconstructions in the Upper cretaceous Haidach Section (Gosau Group; Northern Calcarous Alps, Austria). *Facies* 36:69–90 doi: 10.1007/BF02536878
- Sanders, D., Pons, J.M. and Caus, E. (2004): Shallow-water limestone clasts in a Campanian deep-water debrite (Krappfeld, Central Alps, Austria): implications for carbonate platform history. *Annalen des Naturhistorischen Museums, Wien*. 106A, 139–165.
- Schlagentweit, F. and Rashidi, K., (2016): Some new and poorly known benthic foraminifera from the late Maastrichtian shallow-water carbonates of the Zagros Zone, SW Iran. *Acta Palaeontologica Romaniae*, 12, 1, 53–70.
- Schlüter, M., Steuber, T. and Parente, M., (2008): Chronostratigraphy of Campanian- Maastrichtian platform carbonates and rudist associations of Salento (Apulia, Italy). *Cretaceous Research*, 29, 100–114. Doi: 10.1016/j.cretres.2007.14.005
- Skelton, P.W. and Gili, E. (2002): Paleoecological classification of rudist morphotypes. In: Sladić-Trifunović M. (ed.): Rudists, Proceedings, First International Conference on Rudists. Beograd, 1988, Union of Geological Societies of Yugoslavia, Memorial Publication, Beograd, 71–86.
- Sohl, N. F. and Kollmann, H. A. 1985: Cretaceous Actaeonellid Gastropods from the Western Hemisphere. Geological Survey Professional Paper, 1304, 1–104, pls. 1–23.
- Solak, C., Taslı, K. and Koç H. (2017): Biostratigraphy and facies analysis of the Upper Cretaceous-Danian? platform carbonate succession in the Kuyucak area, western Central Taurides, S Turkey. *Cretaceous Research*, 79, 43–63. doi:10.1016/j.cretres.2017.06.019
- Sremac, J., Bošnjak Makovec, M., Vrsaljko, D., Karaica, B., Tripalo, K., Fio Firi, K., Majstorović Bušić, A. and Marjanac, T. (2016): Reefs and bioaccumulations in the Miocene deposits of the North Croatian Basin – Amazing diversity yet to be described. *Rudarsko-geološko-naftni zbornik*. 31, 1, 19–29. doi: 10.17794/rgn.2016.1.2
- Sremac, J., Tripalo, K., Repac, M., Bošnjak, M., Vrsaljko, D., Marjanac, T., Moro, A., Lužar-Oberiter, B., Fio Firi, K. and Aščić, Š. (2018): Middle Miocene drowned ramp in the vicinity of Marija Bistrica (northern Croatia). *Rudarsko-geološko-naftni zbornik*. 42, (in press). doi: 10.17794/rgn.2018.4.3
- Steuber, T., Korbar, T., Jelaska, V. and Gušić, I. (2005): Strontium-isotope stratigraphy of Upper Cretaceous platform carbonates of the island of Brač (Adriatic Sea, Croatia): Implications for global correlation of platform evolution and biostratigraphy. *Cretaceous Research*. 26, 5, 741–756. doi: 10.1016/j.cretres.2005.04.004.
- Steuber, T. and Schüter, M. (2012): Strontium-isotope stratigraphy of Upper Cretaceous rudist bivalves: Biozones, evolutionary patterns and sea-level change calibrated to numerical ages. *Earth-Science Reviews*. 114, 42–60. doi: 10.1016/j.earscirev.2012.04.004
- Velić, I. (2007): Stratigraphy and Palaeobiogeography of Mesozoic Benthic Foraminifera of the Karst Dinarids (SE Europe). *Geologia Croatica*. 60, 1, 1–113. doi: 10.4154/GC.2007.01
- Vlahović, I., Tišljari, J., Velić, I. and Matičec, D. (2005): Evolution of the Adriatic Carbonate Platform: Palaeogeography, main events and depositional dynamics. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 220, 333–360. doi: 10.1016/j.palaeo.2005.01.001
- Willingshofer, E., Neubauer, F. and Cloetingh, S. (1999): The significance of Gosau-type basins for the Late Cretaceous history of the Alpine-Carpathian belt. *Physics and Chemistry of the Earth (A)*, 24, 8, 687–695.
- Wilmsen, M., Berensmeier, M., Fürsch, F.T., Majidifard, M.R. and Schlagintweit, F. (2018): A Late Cretaceous epeiric carbonate platform: the Haftoman Formation of central Iran. *Facies*, 64, doi: 10.1007/s10347-018-0523-6.

SAŽETAK

Mikrofacijesna obilježja karbonatnih valutica kampana iz prostora Slovenj Gradeca (Slovenija): posljedice za određivanje paleoniše *Fleuryana adriatica* De Castro, Drobne i Gušić te širenje njihova biostratigrafskoga raspona u prostoru Tetisa

U potoku blizu izdanka s kampanskim naslagama u području Slovenj Gradeca uzet je uzorak dimenzija krupnoga šljunka s fosilnim ostacima puževa roda *Trochachtaeon*. Zajedno s njima određene su bentičke foraminifere *Fleuryana adriatica* i *Cuneolina ketini* unutar potplimskoga okoliša s aglutaniniranim i porculanskim foraminiferama koje dosada nisu nađene u ovome dijelu gornjokrednoga oceana Tetis. Stratigrafski podatci iz okolnih gornjokrednih naslaga, zajedno s određenom mikrofossilnom zajednicom iz spomenutoga uzorka krupnoga šljunka, upućuju na kronostratigrafsku starost koja odgovara gornjemu dijelu donjega kampana, što podrazumijeva širi stratigrafski raspon vrsta *F. adriatica* i *C. ketini*. Paleoniša, u kojoj je *F. adriatica* obitavala najpliči je dio potplimskoga okoliša. Obitavala je na relativno tvrđem substratu koji je rezultat lateralno strmijega paleoreljefa. Izostanak vrste *F. adriatica* unutar muljnih karbonatnih naslaga potplimskoga okoliša može biti zbog relativno veće brzine sedimentacije, te u paleookolišima relativno povišene energije koje karakteriziraju grejnston-pekston varijeteti podloge.

Ključne riječi:

gornja kreda, *Fleuryana adriatica*, *Cuneolina ketini*, bentičke foraminifere, paleoniša, karbonatne naslage

Authors' contributions

Alan Moro provided the micropaleontological analysis, palaeoenvironmental interpretations and presentation of the results. **Ivo Velić** provided the micropaleontological analyses of *F. adriatica* and *C. ketini*. **Vasja Mikuž** provided the gastropod paleoecology. **Aleksander Horvat** performed the field work, contributing with the geology of the Gosau deposits.